


Nina R. Ingle<sup>1,3</sup>, Jodi L. Sedlock<sup>2,3</sup>, Lawrence R. Heaney<sup>3</sup> <sup>1</sup>Cornell University, <sup>2</sup>University of Illinois at Chicago, and <sup>3</sup>The Field Museum

Illustrations by J.L. Sedlock. Photos by: J.L. Sedlock, P.D. Heideman, N.R. Ingle, L.R. Heaney, R.L. Baylomo, E.K. Walker, A.C. Diesmos. Produced by: R.B. Foster, N.R. Ingle, M.R. Metz, with support from the Andrew Mellon Foundation, the Chicago Zoological Society, Bat Conservation International, Cornell University, the MacArthur Foundation, and the Brown Fund of The Field Museum. © N.R. Ingle, J.L. Sedlock, L.R. Heaney; Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605, USA. [RRC@fmnh.org] Rapid Color Guide #49 version 1.2



**Bats Recorded from Mindanao**  
 A total of 53 species of bats are known from Mindanao; additional species are probably present. The length of the forearm in adults, often useful for identification, is provided in parentheses. This measurement can be taken with a ruler, but calipers provide more accuracy.

**EMBALLONURIDAE**  
*Emballonura alecto* (44-49 mm)  
*Saccolaimus saccolaimus* (66-71 mm)  
*Taphozous melanopogon* (61-65 mm)

**MOLOSSIDAE**  
*Chaerephon plicata* (40-43 mm)  
*Cheiromeles parvidens* (73-78 mm)  
*Mops sarasinorum* (39-45 mm)

**PTEROPODIDAE**  
*Acerodon jubatus* (165-215 mm)  
*Alionycteris paucidentata* (45-50 mm)  
*Cynopterus brachyotis* (58-68 mm)  
*Dyacopterus spadiceus* (92 mm)  
*Eonycteris robusta* (67-82 mm)  
*Eonycteris spelaea* (67-80 mm)  
*Haplonycteris fischeri* (46-53 mm)  
*Harpionycteris whiteheadi* (80-91 mm)  
*Macroglossus minimus* (41-45 mm)  
*Megaerops wetmorei* (45-52 mm)  
*Ptenochirus jadori* (76-90 mm)  
*Ptenochirus minor* (60-78 mm)  
*Pteropus hypomelanus* (136-149 mm)  
*Pteropus pumilus* (94-113 mm)  
*Pteropus speciosus* (118-133 mm)  
*Pteropus vampyrus* (179-204 mm)  
*Rousettus amplexicaudatus* (80-92 mm)

**RHINOLOPHIDAE**  
*Coelops hirsutus* (34 mm)  
*Hipposideros ater* (38-43 mm)  
*Hipposideros cervinus* (44-50 mm)  
*Hipposideros coronatus* (47 mm)  
*Hipposideros diadema* (77-89 mm)  
*Hipposideros obscurus* (42-48 mm)  
*Rhinolophus arcuatus* -small (43-46 mm)  
*Rhinolophus arcuatus* -large (47-50 mm)  
*Rhinolophus inops* (49-55 mm)  
*Rhinolophus macrotis* (43-44 mm)  
*Rhinolophus philippinensis* (51-57 mm)  
*Rhinolophus rufus* (68-73 mm)  
*Rhinolophus subrufus* (51-56 mm)  
*Rhinolophus virgo* (38-44 mm)

**MEGADERMATIDAE**  
*Megaderma spasma* (57-63 mm)

**VESPERTILIONIDAE**  
*Kerivoula hardwickii* (33-35 mm)  
*Kerivoula pellucida* (35 mm)  
*Kerivoula whiteheadi* (30-31 mm)  
*Miniopterus australis* (34-39 mm)  
*Miniopterus schreibersi* (42-46 mm)  
*Miniopterus tristis* (51-55 mm)  
*Murina cyclotis* (36-39 mm)  
*Myotis horsfieldii* (35-38 mm)  
*Myotis macrotarsus* (44-49 mm)  
*Myotis muricola* (30-34 mm)  
*Philetor brachypterus* (34-39 mm)  
*Pipistrellus javanicus* (32-36 mm)  
*Pipistrellus petersi* (38 mm)  
*Pipistrellus stenopterus* (38-40 mm)  
*Pipistrellus tenuis* (30-32 mm)  
*Scotophilus kuhlii* (47-53 mm)